

=>


Uploading C:\Program Files\Stnexp\Queries\10038760d.str

L4 STRUCTURE UPLOADED

=> d 14

L4 HAS NO ANSWERS

L4 STR



G1 H, SH, NH2, X

Structure attributes must be viewed using STN Express query preparation.

=> s 14 sam sss

SAMPLE SEARCH INITIATED 20:50:27 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 13 TO ITERATE

100.0% PROCESSED 13 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 44 TO 476

PROJECTED ANSWERS: 0 TO 0

L5 0 SEA SSS SAM L4

=> s 14 full sss

FULL SEARCH INITIATED 20:50:35 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 225 TO ITERATE

100.0% PROCESSED 225 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.01

L6 1 SEA SSS FUL L4

=> d 16


L6 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2004 ACS on STN

RN 145383-55-7 REGISTRY

CN D-erythro-Pentitol, 1,4-anhydro-2-deoxy-1-C-4-pyrimidinyl- (9CI) (CA INDEX NAME)

FS STEREOSEARCH
 MF C9 H12 N2 O3
 SR CA
 LC STN Files: CA, CAPLUS
 DT.CA CAPLUS document type: Journal
 RL.NP Roles from non-patents: PREP (Preparation)

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> file caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	157.19	160.01

FILE 'CAPLUS' ENTERED AT 20:50:57 ON 23 NOV 2004
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.


FILE COVERS 1907 - 23 Nov 2004 VOL 141 ISS 22
 FILE LAST UPDATED: 22 Nov 2004 (20041122/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l6
 L7 2 L6
 => d fbib abs ti 1-2 l7

L7 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
 AN 1995:195649 CAPLUS
 DN 122:315016
 TI Synthesis of C-nucleosides via radical coupling reaction
 AU Togo, Hideo; Ishigami, Sachiko; Fujii, Misa; Ikuma, Toshihiro; Yokoyama,

Masataka
 CS Fac. Sci., Chiba Univ., Chiba, 263, Japan
 SO Journal of the Chemical Society, Perkin Transactions 1: Organic and
 Bio-Organic Chemistry (1994), (20), 2931-42
 CODEN: JCPRB4; ISSN: 0300-922X
 PB Royal Society of Chemistry
 DT Journal
 LA English
 GI



AB Photolysis of O-acyl derivs. of N-hydroxy-2-thiopyridone, prepared from tetrahydrofuran-2-carboxylic acid, D-ribofuranosylmethanoic acid, and D-ribopyranosylmethanoic acid, gave the corresponding C-nucleoside derivs., e.g. I (B = R, R1), in the presence of heteroarom. compds. via radical pathways. The essential step in this method is a radical coupling reaction of D-ribofuranosyl radical or D-ribopyranosyl radical and some heteroarom. bases. This is a new method for the preparation of C-nucleosides using sugar carboxylic acids.
 TI Synthesis of C-nucleosides via radical coupling reaction

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS on STN
 AN 1993:60015 CAPLUS
 DN 118:60015
 TI A facile preparative method of C-nucleosides
 AU Togo, Hideo; Ishigami, Sachiko; Yokoyama, Masataka
 CS Fac. Sci., Chiba Univ., Chiba, 263, Japan
 SO Chemistry Letters (1992), (9), 1673-6
 CODEN: CMLTAG; ISSN: 0366-7022
 DT Journal
 LA English
 OS CASREACT 118:60015
 AB Facile and general preparative method of C-nucleosides has been achieved via 4 steps starting from 2-deoxy-D-ribose. The key step in this method is the use of radical coupling reaction of 2-deoxy-D-ribofuranosyl radical derivative and some heteroarom. bases.
 TI A facile preparative method of C-nucleosides